

Analysis of Ozone Pollution Characteristics and Impact Factors in the City of Haikou(2016 to 2020)

Cai, J.Z.¹ Wang, S.H.² Hu, J.X.^{3*}

1. Ecological Environmental Protection Administrative Law Enforcement Detachment of Haikou Comprehensive Administrative Law Enforcement Bureau, Haikou, Hainan, China, 570000;
2. Haikou Municipal Ecology and Environment Bureau, Haikou, Hainan, China, 570000;
3. China National Environmental Monitoring Centre, Beijing, China, 100012;

Abstract: To study the characteristics of ozone pollution changes in tropical island cities and their relationship with meteorological factors, daily and hourly air quality automatic monitoring data and meteorological observations in Haikou City from 2016 to 2020 were selected for analysis. The results show that the peak ozone concentration period of Haikou City is significantly different from that of inland cities, mainly in the autumn and winter seasons (from October to December). Regional climate differences are the main reason for the time distribution. The diurnal variation presents an obvious single-peak feature, with a nadir at 8:00 and a peak at 14:00-16:00. In recent years, the range of ozone pollution in urban areas has gradually expanded, and its degree has worsen; ozone concentration is positively correlated with CO concentration and wind speed, and negatively correlated with humidity. The main meteorological factors that affect the change of ozone concentration in Haikou City in autumn and winter are not air pressure and temperature.

Keywords: Haikou City; variation characteristic; precursor; correlation analysis; meteorological factor

DOI: <https://doi.org/10.3974/geodp.2022.03.16>

CSTR: <https://cstr.science.org.cn/CSTR:20146.14.2022.03.16>

Dataset Availability Statement:

The dataset supporting this paper was published and is accessible through the *Digital Journal of Global Change Data Repository* at:<https://doi.org/10.3974/geodb.2022.05.04.V1>. or <https://cstr.science.org.cn/CSTR:20146.11.2022.05.04.V1>.

1 Introduction

In recent years, O₃ has become one of the most critical pollutants affecting urban air quality in China, and O₃ pollution in cities and regions is on the rise^[1]. According to the 2019 China Ecological Environment Bulletin, in 337 cities at prefectural level and above in China, the average annual concentrations of PM_{2.5}, PM₁₀, O₃, SO₂, NO₂ and CO were 36 μg/m³, 63 μg/m³, 148 μg/m³, 11 μg/m³, 27 μg/m³ and 1.4 mg/m³, respectively. Compared with 2018, O₃ concentration has increased, and the proportion of days exceeding the standard is also on the

Received: 10-06-2022; **Accepted:** 11-08-2022; **Published:** 25-09-2022

Foundation: Haikou Ecological and Environmental Bureau (HXSJ-CG-2021102)

***Corresponding Author:** HU Jiaxin, China Environmental Monitoring Station, hujx@cnemc.cn

Data Citation: [1] Cai, J. Z., Wang, S. H., Hu, J. X. Analysis of ozone pollution characteristics and impact factors in the city of Haikou (2016 to 2020) [J]. *Journal of Global Change Data & Discovery*, 2022, 6(3): 452-462. <https://doi.org/10.3974/geodp.2022.03.16>. <https://cstr.science.org.cn/CSTR:20146.14.2022.03.16>.

[2] Cai, J. Z., Wang, S. H., Hu, J. X.. Dataset of Ozone Pollution Characteristics and Impact Factors (2016-2020)[J/DB/OL]. *Digital Journal of Global Change Data Repository*, 2022. <https://doi.org/10.3974/geodb.2022.05.04.V1>. or <https://cstr.science.org.cn/CSTR:20146.11.2022.05.04.V1>.

rise. O₃ pollution has become increasingly prominent^[2]. At present, with the rapid economic development and the continuous expansion of the urban scale, O₃ pollution has become an important environmental problem that urgently needs to be solved in the process of urban development in China and is one of the hot research topics in the field of atmospheric chemistry and environmental science. It is generally believed that the changing trend of O₃ pollution is subject to the joint action of the emission of precursor substances, meteorological conditions, and chemical reactions and that there is a significant correlation between the change of O₃ concentration and meteorological conditions, with obvious seasonal variation characteristics^[3-4]. However, domestic analysis and research on O₃ pollution are mainly concentrated in some key cities and economically developed areas. In recent years, the economic development and urban expansion of Hainan Province have led to a worsening air quality of Hainan Island, and the situation of O₃ pollution has become increasingly severe^[5-6]. Haikou is a tropical island city with unique natural conditions, its O₃ pollution characteristics and meteorological impact have certain typicality. Therefore, based on the concentration data of O₃, CO, and NO₂ from four state-controlled monitoring sites in Haikou City during the 13th Five-Year Plan (2016-2020), this paper analyzes the overall characteristics of O₃ pollution and the potential relationship between O₃ and precursor substances. In addition, the monthly and daily variation characteristics of O₃ concentration at the state-controlled monitoring stations were analyzed. Through studying the relationship between O₃ and meteorological factors, the weather types that are prone to trigger O₃ pollution were discussed, providing a reference for O₃ pollution warning and prevention in Haikou City.

2 Metadata of Dataset

The metadata of “Analysis of ozone pollution characteristics and impact factors in the city of Haikou(2016 to 2020)”^[7] dataset is summarized in Table 1. It includes the dataset’s full name, short name, authors, year of the dataset, temporal resolution, spatial resolution, data format, data size, data files, data publisher, data sharing policy, etc.

3 Methods

3.1 Primary Data

The O₃, CO, NO₂, and corresponding meteorological data (air pressure, air temperature, humidity, and wind speed) from 2016 to 2020 used in this research are respectively from 4 state-controlled automatic air quality monitoring stations in Haikou (distribution of stations is shown in Figure 1) and information center of Hainan Meteorological Bureau. Four state-controlled sites respectively: Haida site, Xiuying site, Haishi site, and Longhua site, The state-controlled stations are representative, comparable, and holistic, and can reflect the status of Haikou City air quality more accurately.

3.2 Algorithm Principle

The evaluation of monitoring results refers to Environmental Air Quality Standard (GB3095-2012) and Technical Specification for Environmental Air Quality Evaluation (TRIAL) (HJ633-2012). The overall pollution characteristics of O₃ in Haikou city from 2016 to 2020 are statistically analyzed based on the average O₃-8h-90PER of 4 stations. CO-95PER and NO₂ are used to analyze the relationship between O₃ concentration and precursors. O₃-8h-90PER and O₃-1h of state-controlled sites are used to study the monthly and daily changes of O₃ pollution, respectively. In addition, Spielman correlation coefficient

method and Pearson correlation coefficient method are used for correlation analysis of the relationship between O₃ and precursors and meteorological factors. Correlation coefficient $0.1 < |r| \leq 0.3$ is a weak correlation, $0.3 < |r| \leq 0.5$ is a moderate correlation, and $|r| > 0.5$ is a strong correlation.

Table 1 Metadata Summary of the Analysis of Ozone Pollution Characteristics and Impact Factors in the City of Haikou(2016 to 2020)

Items	Description
Dataset full name	Dataset of Ozone Pollution Characteristics and Impact Factors (2016-2020)
Dataset short name	Ozone Pollution Characteristics in the City of Haikou
Authors	CAI Jinzi, Ecological Environmental Protection Administrative Law Enforcement Detachment of Haikou Comprehensive Administrative Law Enforcement Bureau, email:caijz@haikou.gov.cn;WANG Shaohu, Haikou Bureau of Ecology and Environment, email:wangshaohui@haikou.gov.cn;HU Jiaxin, China Environmental Monitoring Station, email: hujx@cnemc.cn;
Geographical region	Haikou City
Year	2016 to 2020
Data format	.shp, .xlsx
Data size	33.9 KB
Data files	(1) geo-location of the sites; (2) monthly concentration data of O ₃ , CO, and NO ₂ in Haikou from 2016 to 2020; (3) monthly and annual O ₃ concentration at four monitoring sites from 2016 to 2020; (4) daily variation data of O ₃ concentration at four monitoring sites from 2016 to 2020; (5) O ₃ concentration, air pressure, air temperature, humidity and wind speed data at four monitoring sites. It consists of 9 data files.
Foundation(s)	Haikou Ecological and Environmental Bureau (HXSJ-CG-2021102)
Data publisher	Global Change Research Data Publishing & Repository, http://www.geodoi.ac.cn
Address	No. 11A, Datun Road, Chaoyang District, Beijing 100101, China
Data sharing policy	<i>Data</i> from the Global Change Research Data Publishing & Repository includes metadata, datasets (in the <i>Digital Journal of Global Change Data Repository</i>), and publications (in the <i>Journal of Global Change Data & Discovery</i>). <i>Data</i> sharing policy include: (1) <i>Data</i> are openly available and can be freely downloaded via the Internet; (2) End users are encouraged to use <i>Data</i> subject to citation; (3) Users, who are by definition also value-added service providers, are welcome to redistribute <i>Data</i> subject to written permission from the GCdataPR Editorial Office and the issuance of a <i>Data</i> redistribution license; and (4) If <i>Data</i> are used to compile new datasets, the ‘‘ten percent principal’’ should be followed such that <i>Data</i> records utilized should not surpass 10% of the new dataset contents, while sources should be clearly noted in suitable places in the new dataset ^[7]
Communication and searchable system	DOI, CSTR, Crossref, DCI, CSCD, CNKI, SciEngine, WDS/ISC, GEOSS




Fig.1 Air quality monitoring station in Haikou City

4 Results and Validation

4.1 Data Composition

The dataset includes (1) geo-location of the sites; (2) monthly concentration data of O₃, CO, and NO₂ in Haikou from 2016 to 2020; (3) monthly and annual O₃ concentration at four monitoring sites from 2016 to 2020; (4) daily variation data of O₃ concentration at four monitoring sites from 2016 to 2020; (5) O₃ concentration, air pressure, air temperature, humidity, and wind speed data at four monitoring sites. The dataset is archived in .shp and .xlsx data formats; it consists of 9 data files.

4.2 Data Products

4.2.1 Overall characteristics of O₃ pollution

Figure 2 shows the monthly changes of O₃-8h-90PER average concentration, CO-95PER, and NO₂ average concentration from 2016 to 2020. It can be seen from Figure 2, that the concentration peaks of O₃ in each year appear from September to December, which is 136, 163, 183, 178, and 156 μg/m³, respectively. The overall trend of O₃ pollution increased year by year from 2016 to 2019 and decreased in 2020. In addition, the peak of O₃ concentration from 2017 to 2019 exceeded the secondary limit concentration stipulated in the standard, which preliminarily indicated that O₃ pollution in Haikou City was more severe in autumn and winter than in summer. On the one hand, this phenomenon is related to the fact that Haikou is rainy in summer, low solar radiation, and wet deposition are not conducive to O₃ generation^[8]. On the other hand, Hainan Province may be affected by exogenous transmission, as it is affected by surface cold high pressure control in autumn and winter, the weather situation is stable, and the low-level northerly wind field is controlled, . Under the joint action of local emission and exogenous transport, O₃ concentration increases^[9]. In addition, Haikou City boasts the relatively high temperature in autumn , belonging to tropical monsoon climate with moderate solar radiation intensity and good photochemical reaction, which is conducive to the O₃ generation and accumulation . After December, cold air invades ,and O₃ concentration decreases, so O₃ concentration reaches the peak from September to December. Compared with other years, the average daily concentration of O₃ in 2016 had the smallest variation range, and its concentration was mainly distributed between 84 and 136 μg/m³. The concentration of O₃ in 2017 had the largest variation range, and its concentration was mainly distributed between 60 and 163 μg/m³. Meanwhile, the concentration of O₃ in 2016 and 2020 were below the secondary limit. The O₃ concentration was generally low.




Fig.2 Monthly variations of O₃-8h and CO and NO₂ from 2016 to 2020 (blue line: the primary limit of 100 μg/m³, red line: the secondary limit of 160 μg/m³)

Figure 2 shows the correlation coefficient between O₃-8h and the concentration of CO and NO₂. It could be seen that the correlation coefficient between O₃ and CO was 0.552, indicating a strong positive correlation and that between O₃ and NO₂ was -0.14, indicating a weak negative correlation (Table 2). From the view of the monthly average concentration distribution of CO and O₃, CO and O₃ showed positive phase changes in most periods from 2016 to 2020. In 2016, both CO and O₃ had their highest concentration values in December. In 2017-2020, when O₃ concentration increased sharply in October and November, the concentration of CO in the corresponding period also showed an obvious increase. From the perspective of O₃ photochemical formation process, CO is constantly oxidized, so in most cases, CO and O₃ are negatively correlated^[10], while Haikou is positively correlated, which is speculated to be related to the pollution degree of air mass^[11]. In addition, with the rapid growth of vehicles in Haikou City, a great deal of automobile exhaust emissions make the amount of CO increase, causing pollution to the air environment. The variation trend of O₃ concentration is affected by many factors such as precursor concentration, meteorological conditions, and chemical reactions. In terms of the monthly mean concentration distribution of NO₂ and O₃, they showed negative phase change in all months except February and March in 2016, negative phase change in most periods from 2017 to 2020, and positive phase change in a short period in summer. In Hainan Province, because of high summer temperature and sufficient sunshine duration, photochemical reactions occur under the influence of strong solar radiation, so O₃ concentration usually reaches the annual maximum value^[12]. However, summer in Hainan Province is the primary flood season with abundant rainfall. The erosion of rain and high relative humidity both inhibit the photochemical reaction to some extent and weaken the influence of NO₂ and O₃, so O₃ concentration does not reach the maximum in summer^[13-14]. In addition, when the concentration of CO and NO₂ reached a relatively stable plateau, O₃ concentration still fluctuated, which reflected the complexity of factors influencing O₃ concentration change in Haikou City. The precursor could not fully represent the occurrence of O₃ pollution, so it was necessary to further analyze the temporal variation characteristics of O₃ concentration and the influence of meteorological factors.

The monthly mean concentration of O₃-8h-90PER for the Haida site, Xiuying site, Haishi site, and Longhua site during 2016-2020 were statistically analyzed, and the outline diagram of the O₃ concentration of each station was drawn with the maximum, minimum, and average values as characteristic values (Figure 3). In our study, the average values of the maximum O₃ concentration at the Haida site, Xiuying site, Haishi site, and Longhua site were 105, 85, 94, 109, 92 μg/m³, and the average values of the minimum O₃ concentration were 53, 36, 31, 36, 31 μg/m³, respectively. The O₃ concentration were 74, 62, 55, 64, and 60 μg/m³ by

Table 2 The correlation coefficient between O₃-8h concentration and precursor concentration in 2016-2020

Category	Correlation Index	P Value
O ₃ -CO	0.552	$p < 0.01$
O ₃ -NO ₂	-0.14	$p > 0.05$




Fig. 3 2016-2020 annual average O₃ concentration of 4 urban testing sites

average analysis of the average values of each site. As can be seen from the above O₃ concentration data, compared with developed areas such as Guangdong and Chengdu Province, the O₃ pollution level in Haikou City had not reached a significantly high pollution level^[14,17,20]. According to the analysis of the O₃ concentration of the four sites, the Haida site was in the high-value area in 2016, while the Xiuying site was in the low-value area. From 2017 to 2020, the high-value area and the low-value area were no longer noticeable, indicating that the area scope of O₃ pollution in the urban area of Haikou was expanding.

4.2.2 Time-varying characteristics of O₃ concentration

4.2.1.1 Monthly variation characteristics of point position

Figure 4 shows the monthly variation of the mean concentration of O₃-8h-90PER at state-controlled monitoring sites from 2016 to 2020. According to the spring (March to May), summer (June to August), autumn (September to November), and winter (December to the next February) classification. It could be seen that the seasonal variation of O₃ was apparent. The peak value of O₃ concentration at the four sites occurred frequently in October-December (autumn and winter), in which the average maximum value of O₃ in 2016 and 2017 appeared in December, the average maximum value of O₃ in 2018 appeared in October, and the average maximum value of O₃ in 2019 and 2020 appeared in November. This is consistent with the time range of the peak O₃ concentration in Haikou City from 2016 to 2020. This is different from cities with severe air pollution in China. For example, O₃ concentration in Guangdong and Shanghai in summer is higher than that in autumn and winter^[14-17]. The high concentration of O₃ in Haikou City in autumn and winter was mainly due to the northeast wind in winter, which was easily affected by inland pollution transport. Besides, the temperature in autumn was not low, and the photochemical reaction conditions were good, which was conducive to the generation of O₃^[5]. In addition, Haikou City has frequent typhoons in October, and the pollutants are not easy to diffuse under the action of external downdraft. Further analysis found that in July 2016 and May 2017 - each site's O₃ concentration was on a decline trend, August 4, 2020 - a downward trend, refer to the corresponding climate data showed that during the period while the temperature was higher, the Haikou City was mainly influenced by the southwest monsoon, clean air from the ocean of air dilute the atmosphere in Haikou City. The regional climate difference was the main reason why the time distribution of O₃ in Haikou was different from that in inland cities. It can also be seen from Fig. 4, that O₃ in January-February 2016 was in a rising stage, while all other years were in a declining stage. O₃ concentration in January-March 2019 was lower




Fig. 4 Monthly variations of O₃ average concentration at national monitoring sites from 2016 to 2020

than that in other years, indicating that it was necessary to deal with O₃ spring pollution in advance. In addition, O₃ from August to October 2016 to 2020 presented a trend of aggravation, with a significantly higher growth rate than in other months, further highlighting the severity of the pollution situation in autumn, indicating that this period was the key period for O₃ pollution prevention and control.

4.2.2.2 Diurnal variation characteristics of point position

Figure 5 shows the diurnal variation of O₃-1h mean concentration at the state-controlled monitoring sites from 2016 to 2020. It could be seen that the variation characteristics of O₃ concentration at the four stations were single-peak type, and the maximum O₃ concentration from 14:00 to 16:00 throughout the day, which was similar to the diurnal variation pattern of O₃ concentration in other cities in China [18-19]. This variation was due to the daily budget mechanism of O₃ and was related to the photochemical reaction rate and atmospheric diffusion capacity. The concentration of O₃ from 00:00 to 8:00 showed a trend of continuous decline and decreased to the lowest value at 8:00, with a trough value. This is because the anthropogenic activities weakened during this period, the emissions of precursor substances decreased, the night temperature was low, and the photochemical reaction was almost zero. O₃ is mainly consumed, and its budget is negative, so it remains in the range of low concentration value. The concentration of O₃ from 8:00 to 16:00 showed a trend of continuous rise and reached the peak concentration in a day. This is because the solar radiation intensity keeps increasing after sunrise, and the temperature gradually rises. In addition, the concentration of O₃ precursor produced by transportation sources, industrial sources, and biological sources increases, and the photochemical reaction rate increases. After 16:00, O₃ concentration gradually returned to the low range; this is mainly due to the solar radiation after 16:00 started to wane, vertical mixing, horizontal divergence strengthened and weakened photochemical reaction, and NO peak emissions during rush hour consumes O₃, the accumulation of the O₃ return to give way to consumption function, with balance negative. Then a concentration decreases continuously, and the cycle continues [20-21]. In addition, the overall level of O₃ daily concentration in 2019 was higher than that in other years, which may be related to the strengthening of urban emissions and meteorological conditions.




Fig. 5 Daily variations of O₃ average concentration at national monitoring sites from 2016 to 2020

4.2.3 Influence of meteorological factors on typical O₃ pollution days

Meteorological factors can affect the formation, transmission, settlement, and dissipation of O₃. Based on the analysis of the above results, it could be known that the high-incidence period of O₃ pollution in Haikou City was autumn and winter (September to December). To better study the relationship between O₃ concentration and meteorological factors, the O₃-8h

concentration data and meteorological factor data of the same period from September to December 2016 to 2020 from four state-controlled monitoring stations were selected for analysis, and the correlation coefficients between O₃-8h and various meteorological factors were calculated (Table 3). The characteristics of meteorological factors that easily trigger O₃ pollution weather in Haikou City were obtained.

Table 3 Correlation coefficient between O₃ and various meteorological factors

Monitoring Site	O ₃ -Pressure		O ₃ -Temperature		O ₃ -Humidity		O ₃ -Wind Speed	
	Correlation Index	P Value	Correlation Index	P Value	Correlation Index	P Value	Correlation Index	P Value
Haida site	0.398	<i>p</i> >0.05	0.417	<i>p</i> >0.05	-0.539	<i>p</i> <0.05	0.630	<i>p</i> <0.01
Haishi site	0.216	<i>p</i> >0.05	0.374	<i>p</i> >0.05	-0.170	<i>p</i> >0.05	0.327	<i>p</i> >0.05
Longhua site	0.244	<i>p</i> >0.05	0.032	<i>p</i> >0.05	-0.425	<i>p</i> >0.05	0.454	<i>p</i> <0.05
Xiuying site	0.553	<i>p</i> <0.01	0.193	<i>p</i> >0.05	-0.622	<i>p</i> <0.01	0.297	<i>p</i> >0.05

4.2.3.1 Influence of air pressure and temperature on O₃ pollution

Fig. 6 shows the changes of ozone and air pressure, and temperature at national monitoring sites. It can be seen that O₃ concentration in Haikou is positively correlated with air pressure and temperature. Quantitative analysis of the relationship between them (Table 2) showed that the correlation coefficient between O₃ concentration and air pressure is positive. However, only the positive correlation at the Xiuying site passed the significance test (*P* < 0.01), and the correlation between O₃ concentration and air pressure was weak at other three stations. The correlation coefficient between O₃ concentration and the air temperature was positive, but the correlation coefficient between O₃ concentration and air temperature at four stations did not pass the significance test (*P* > 0.05). Haikou City is located at the tropical edge of low latitude and has a tropical maritime monsoon climate. It is cool in autumn, and the temperature is lower than that in spring and summer. In winter, due to the southward extension of the continental cold air mass, there often invades cold airflow, which indicates that the main meteorological factors affecting the change of O₃ concentration in Haikou City from September to December are not atmospheric pressure and temperature.




Fig. 6 Changes of ozone and air pressure and temperature at national monitoring sites

4.2.3.2 Influence of humidity and wind speed on O₃ pollution

Figure 7 shows the change of O₃ concentration, humidity, and wind speed at the state-controlled monitoring site. It could be seen that O₃ concentration was negatively correlated with humidity and positively correlated with wind speed. Quantitative analysis of the relationship between O₃ concentration, humidity, and wind speed (Table 2) manifested that the correlation coefficient between O₃ and humidity at all monitoring stations was negative. There was a significant negative correlation between O₃ concentration and humidity at the Haida site ($p < 0.05$), while there was a strong negative correlation between O₃ concentration and humidity at the Xiuying site ($p < 0.01$). The correlation coefficients between O₃ and wind speed at all monitoring stations were positive. There was a significant positive correlation between O₃ concentration and wind speed at the Longhua site ($p < 0.05$) and a strong positive correlation between O₃ concentration and wind speed at the Haida site ($p < 0.01$). It indicated that from September to December, the change of O₃ concentration in Haikou City was more likely to be negatively affected by humidity and positively affected by wind speed.




Fig. 7 Changes of ozone and wind speed and humidity at national monitoring sites

Haikou is prone to heavy rain in autumn and sometimes cloudy in winter, so the O₃ concentration and humidity trend in Haikou is opposite from September to December. Wind speed can reflect pollutant transport efficiency and removal efficiency^[22]. The increase of wind speed has dual effects on O₃ concentration, which occurs simultaneously. Firstly, the air mass transport power increases, the height of the atmospheric boundary layer increases and the vertical momentum transport is enhanced, all that are conducive to the transmission of O₃ from the upper region to the ground. When accumulation remains dominant in local areas, the concentration of O₃ will increase. Secondly, the horizontal diffusion effect is enhanced. When it is dominant, the increase of wind speed accelerates O₃ dilution^[23]. When the wind speed decreases, its scavenging effect on O₃ is limited, and the cumulative effect is dominant, increasing the concentration of O₃^[24]. It can be seen from Figure 8 the wind speed monitored by the four state-controlled sites from September to December 2016-2020 ranges from 0.1 to 3.8 m s⁻¹. The wind speed was low, and the vertical downward transport

effect of O₃ was stronger than the horizontal diffusion effect, which was conducive to the accumulation of O₃. Therefore, there is a positive correlation between O₃ concentration and wind speed in Haikou from September to December. On the basis of strengthening the existing pollution prevention and control in Haikou City, strengthening joint prevention and control with upper-level regions is the key to effectively control O₃ secondary pollution.

To sum up, in autumn and winter (September to December) when O₃ pollution occurs frequently, the influence of meteorological factors on O₃ concentration is in the following order --humidity > wind speed > air pressure > temperature. It indicated that humidity and wind speed were the main meteorological factors affecting O₃ pollution in Haikou City in autumn and winter. On the whole, low humidity and low wind speed were prone to O₃ pollution. The change of O₃ concentration was affected by many factors, such as precursors, atmospheric fine particles, and meteorological factors, and the changing process was a complex and comprehensive interaction process. Therefore, the influence of meteorological factors on O₃ concentration was inevitably affected by other non-meteorological factors and the interaction between meteorological factors. Consequently, in the study of O₃ pollution characteristics and meteorological impact, to improve the accuracy of O₃ pollution prevention and control in Haikou City, we must consider the role of various possible factors, adjust pollution prevention and control strategies timely, and strengthen the cooperative control of multiple regions.

5 Discussion and Conclusion

(1) From 2016 to 2020, the peak occurrence period of O₃ concentration in Haikou City is significantly different from other inland cities, mainly appeared in autumn and winter, from October to December. Regional climate difference is the main reason why the time distribution of O₃ in Haikou is different from that in inland cities.

(2) The diurnal variation of O₃ concentration presents a prominent single-peak characteristic, with a trough value at 8:00 and a peak value from 14:00 to 16:00. According to the characteristic value of O₃ concentration, the scope of O₃ pollution in urban areas has gradually expanded and worsened in recent years.

(3) O₃ concentration is positively correlated with CO concentration and wind speed, negatively correlated with humidity. Air pressure and temperature are not the main meteorological factors affecting O₃ concentration change in Haikou City in autumn and winter .

(4) In line with the feature of O₃ pollution in Haikou city, a series of measures have been taken, such as stimulating the purchase of new energy vehicles, encouraging the district government and relevant departments to establish a responsibility system, industrial enterprises to implement a staggered rush hour production plan and motor vehicle peak shifting travel, improving pollution prevention and control measures of the third service catering enterprises , strengthening the environmental monitoring and supervising capacity, promoting joint prevention and control with upper-level regions. O₃ pollution in Haikou City is expected to be improved.

Author Contributions

Cai, J.Z designed the algorithms of the dataset. Wang, S.H. contributed to the data collecting and processing. Hu, J.X wrote the data paper.

Acknowledgements

This manuscript is funded by “Environmental Protection Special Fund Project: Haikou Fine Particulate Matter and Ozone Pollution Collaborative Prevention and Control Station Tracking Service Project (HXSJ-CG-2021102)”.

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] Wang, T., Xue, L.K., Brimblecombe, P., et al. Ozone pollution in China: a review of concentrations, meteorological influences, chemical precursors, and effects[J]. *Science of the Total Environment*, 2017, 575: 1582–1596.
- [2] Ministry of Ecology and Environment. 2019 China's Ecological and Environmental Status Bulletin [EB/OL]. (2020-06-02)[2021-5-7].<http://www.mee.gov.cn/hjzl/sthjzk/zghjzkgb/202006/P020200602509464172096.pdf>.
- [3] Yang, L., Luo, H., Yuan, Z., et al. Quantitative impacts of meteorology and precursor emission changes on the long-term trend of ambient ozone over the pearl river delta, China, and implications for ozone control strategy[J]. *Atmospheric Chemistry and Physics*, 2019, 19: 12901–12916.
- [4] Jacob, D.J., Winner, D.A. Effect of Climate Change on Air Quality[J]. *Atmospheric Environment*, 2009, 43(1): 51–63.
- [5] Xu, W.S., Xing, Q., Meng, X.X., et al. Characteristics of ozone pollution in Haikou city [J]. *China Environmental Monitoring*, 2017, 33(4): 186–193. (in Chinese)
- [6] Fu, C.B., Dan, L., TONG, J.H., et al. Variation characteristics of ozone concentration in Haikou urban area during 2013-2018 [J]. *China Environmental Monitoring*, 2020, 36(05): 38–46. (in Chinese)
- [7] CAI Jinzi, WANG Shaohui, HU Jiabin*, et al. Dataset of Ozone Pollution Characteristics and Impact Factors (2016-2020)[J/DB/OL]. Digital Journal of Global Change Data Repository, 2022. <https://doi.org/10.3974/geodb.2022.05.04.V1>.
- [8] Awang, N.R., Ramli, N.A., Yahaya, A.S., et al. Multivariate methods to predict ground-level ozone during the daytime, nighttime, and critical conversion time in urban areas[J]. *Atmospheric Pollution Research*, 2015, 6(5): 726–734.
- [9] Fu, C.B., Chen, Y.L., Dan, L., et al. Temporal and spatial variation of atmospheric NO₂ in Hainan island in recent 10 Years and analysis of pollutant sources [J]. *Environmental Science*, 2015, 37(9): 18–24. (in Chinese)
- [10] Cao, T.W., Wu, K., Kang, P., et al. Characteristics and influencing factors of ozone pollution in Chengdu-Chongqing urban agglomeration [J]. *Chinese Journal of Environmental Science*, 2017, 38(4): 1275–1284. (in Chinese)
- [11] Yang, Y.X., Chen, N.H., Hu, B.Y., et al. Characteristics and influencing factors of ozone pollution in clean islands on the west coast of Taiwan strait [J]. *Environmental Chemistry*, 2020, 39(07): 1733–1743. (in Chinese)
- [12] Fu, C.B., Xu, W.S., Dan, L., et al. Effects of precursors and meteorological factors on ozone pollution in Hainan Province [J]. *Environmental Science and Technology*, 2020, 43(07): 45–50. (in Chinese)
- [13] Dong, H., Cheng, L., Wang, H.Y., et al. Analysis of ozone pollution characteristics and meteorological influencing factors in Anhui Province [J]. *China Environmental Monitoring*, 2021, 37(01): 58–68. (in Chinese)
- [14] Xu, K., Liu, Z.H., He, M.Q., et al. Meteorological characteristics of near-surface ozone pollution in Chengdu City in summer [J]. *China Environmental Monitoring*, 2018, 34(5): 41–50. (in Chinese)
- [15] Liu, Z., Zhu, Y.F., Guo, W.K., et al. Ozone and secondary organic aerosol generation potential of VOCs emission from fossil fuel combustion sources in Lanzhou city [J]. *Environmental science*, 2019, 40(5): 2069–2077. (in Chinese)
- [16] Zhao, C.H., Geng, F.H., Ma, C.Y., et al. Study on aerosol characteristics of Photochemical Pollution in Shanghai area [J]. *China Environmental Science*, 2015, 35(2): 356–363. (in Chinese)
- [17] Shen, J., Huang, X.B., Wang, Y., et al. Characteristics and sources of ozone pollution in Guangdong Province [J]. *Chinese Journal of Environmental Science*, 2017, 37(12): 4449–4457. (in Chinese)
- [18] Pan, B.F., Cheng, L.J., Wang, J.G., et al. Characteristics and source analysis of ozone pollution in Beijing-Tianjin-Hebei region [J]. *China Environmental Monitoring*, 2016, 32(5): 17–23. (in Chinese)
- [19] Yu, S.J., Yin, S.S., Zhang, R.Q., et al. Analysis of ozone pollution characteristics and meteorological factors in Zhengzhou city [J]. *China Environmental Monitoring*, 2017, 33(4): 140–149. (in Chinese)
- [20] Wu, K., Kang, P., Wang, Z.S., et al. Analysis of ozone pollution characteristics and meteorological causes in Chengdu City [J]. *Journal of Environmental Science*, 2017, 37(11): 4241–4252. (in Chinese)
- [21] Song, X.Y., Gao, L.Z., Luo, D., et al. Analysis of ozone Pollution characteristics and Meteorological Impact in Yunnan Province [J]. *China Environmental Monitoring*, 2020, 36(04): 16–28. (in Chinese)
- [22] Yan, R.S., Chen, M.D., Gao, Q.X., et al. Distribution characteristics and influencing factors of typical ozone pollution in Beijing in summer [J]. *Environmental Science Research*, 2013(1): 47–53. (in Chinese)
- [23] Xie, W. J., Wang, S. F., He, L. Y., et al. Characteristics of ozone pollution and its relationship with meteorological elements in Guigang city during 2015-2019 [J]. *Meteorological Research and Application*, 2021, 42(1): 58–62.
- [24] Li, S.J., Li, H., Chen, M., et al. Influence of meteorological factors on ozone and its precursors in the atmosphere of Southwest urban area of Xi'an [J]. *Journal of Meteorology and Environment*, 2018, 34(4): 59–67. (in Chinese)